

# QUANTUM MATRIX BALL: THE WEIGHTED BERGMAN KERNELS

*D. Shklyarov*

*S. Sinel'shchikov*

*L. Vaksman*

Institute for Low Temperature Physics & Engineering  
National Academy of Sciences of Ukraine

The Cartan domains are among the important subjects in many problems of representation theory and mathematical physics [1, 4]. The methods of quantum groups theory [2] were used in [7] to produce  $q$ -analogues of Cartan domains, in particular,  $q$ -analogues of balls in the spaces of complex matrices.

The point of this work is to consider those quantum matrix balls and the associated Hilbert spaces of 'functions'. As a main result, we present an explicit formula for the weighted Bergman kernel.

It is implicit everywhere in the sequel that  $q \in (0, 1)$ ,  $m, n \in \mathbb{N}$ , and  $m \leq n$ .

We need  $q$ -analogues for  $*$ -algebras  $\text{Pol}(\text{Mat}_{mn})$  of polynomials on the space  $\text{Mat}_{mn}$  of complex matrices and a  $*$ -algebra  $D(\mathbb{U})$  of smooth finite functions in the matrix ball  $\mathbb{U} = \{A \in \text{Mat}_{mn} \mid \|A\| < 1\}$ . We start with forming a  $q$ -analogue for the algebra  $\text{Fun}(\mathbb{U}) = \text{Pol}(\text{Mat}_{mn}) + D(\mathbb{U})$ .

The  $*$ -algebra  $\text{Fun}(\mathbb{U})_q$  is given by its generators  $f_0, z_a^\alpha$ ,  $a = 1, 2, \dots, n$ ,  $\alpha = 1, 2, \dots, m$ , and the relations

$$\begin{cases} z_a^\alpha z_b^\beta = q z_b^\beta z_a^\alpha, & a = b \quad \& \quad \alpha < \beta \quad \text{or} \quad a < b \quad \& \quad \alpha = \beta \\ z_a^\alpha z_b^\beta = z_b^\beta z_a^\alpha, & \alpha < \beta \quad \& \quad a > b \\ z_a^\alpha z_b^\beta - z_b^\beta z_a^\alpha = (q - q^{-1}) z_a^\beta z_b^\alpha, & \alpha < \beta \quad \& \quad a < b \end{cases} \quad (1)$$

$$(z_b^\beta)^* \cdot z_a^\alpha = q^2 \sum_{a', b'=1}^n \sum_{\alpha', \beta'=1}^m R(b, a, b', a') R(\beta, \alpha, \beta', \alpha') \cdot z_{a'}^{\alpha'} \cdot (z_{b'}^{\beta'})^* + (1 - q^2) \delta_{ab} \delta^{\alpha\beta}, \quad (2)$$

$$(z_a^\alpha)^* f_0 = f_0 z_a^\alpha = 0, \quad (3)$$

$$f_0 = f_0^* = f_0^2.$$

Here  $a, b = 1, 2, \dots, n$ ,  $\alpha, \beta = 1, 2, \dots, m$ ,

$$R(i, j, i', j') = \begin{cases} q^{-1} & , \quad i \neq j \quad \& \quad i = i' \quad \& \quad j = j' \\ 1 & , \quad i = j = i' = j' \\ -(q^{-2} - 1) & , \quad i = j \quad \& \quad i' = j' \quad \& \quad j' > j \\ 0 & , \quad \text{otherwise} \end{cases}.$$

In this setting, the  $*$ -subalgebra  $\text{Pol}(\text{Mat}_{mn})_q \subset \text{Fun}(\mathbb{U})_q$  generated by  $z_a^\alpha$ ,  $a = 1, 2, \dots, n$ ,  $\alpha = 1, 2, \dots, m$  is a  $q$ -analogue of the  $*$ -algebra  $\text{Pol}(\text{Mat}_{mn})$ , and the bilateral ideal  $D(\mathbb{U})_q =$

$\text{Pol}(\text{Mat}_{mn})_q f_0 \text{Pol}(\text{Mat}_{mn})_q$  is a  $q$ -analogue of the  $*$ -algebra  $D(\mathbb{U})$ . (The element  $f_0$  works here as a delta-function, as one can see from (3)).

To motivate our subsequent constructions, observe that (see [6]) the  $*$ -algebra  $D(\mathbb{U})_q$  is a  $U_q \mathfrak{sl}_{nm}$ -module algebra [2]. Remind the explicit formula for invariant integral from [6].

Consider the representation  $T$  of  $\text{Fun}(\mathbb{U})_q$  in the space  $\mathcal{H} = \text{Fun}(\mathbb{U})_q f_0 = \text{Pol}(\text{Mat}_{mn})_q f_0$ :

$$T(f)\psi = f\psi, \quad f \in \text{Fun}(\mathbb{U})_q, \quad \psi \in \mathcal{H}.$$

There exists a unique positive scalar product in  $\mathcal{H}$  such that  $(f_0, f_0) = 1$ , and

$$(T(f)\psi_1, \psi_2) = (\psi_1, T(f^*)\psi_2), \quad f \in \text{Fun}(\mathbb{U})_q, \quad \psi_1, \psi_2 \in \mathcal{H}.$$

One can prove that the  $*$ -algebra  $\text{Pol}(\text{Mat}_{mn})_q$  admits a unique up to unitary equivalence *faithful* irreducible  $*$ -representation by *bounded* operators in a Hilbert space. This  $*$ -representation can be produced via extending the operators  $T(f)$ ,  $f \in \text{Pol}(\text{Mat}_{mn})_q$ , onto the completion of the pre-Hilbert space  $\mathcal{H}$ .

The invariant integral is of the form (see [6]):

$$\int_{\mathbb{U}_q} f d\nu = \text{tr}(T(f)q^{-2\Gamma(\check{\rho})}), \quad f \in D(\mathbb{U})_q, \quad (4)$$

with  $\Gamma : \mathfrak{h} \rightarrow \text{End}(\mathcal{H})$  being a subrepresentation of the natural representation of the Cartan subalgebra  $\mathfrak{h} \subset \mathfrak{sl}_N$  in  $\text{Fun}(\mathbb{U})_q$ , and  $\check{\rho} \in \mathfrak{h}$  the element of this Cartan subalgebra determined by the half sum of positive roots  $\rho$  under the standard pairing of  $\mathfrak{h}$  and  $\mathfrak{h}^*$ . (To see that this integral is well defined, observe that the operators  $T(f)$ ,  $f \in D(\mathbb{U})_q$ , are finite dimensional, and  $\mathcal{H}$  is decomposable into a sum of weight subspaces associated to non-negative weights.)

Our immediate intention is to produce  $q$ -analogues of weighted Bergman spaces. In the case  $q = 1$  one has

$$\det(1 - \mathbf{z}\mathbf{z}^*) = 1 + \sum_{k=1}^m (-1)^k \mathbf{z}^{\wedge k} \mathbf{z}^{*\wedge k}, \quad (5)$$

with  $\mathbf{z}^{\wedge k}$ ,  $\mathbf{z}^{*\wedge k}$  being the "exterior powers" of the matrices  $\mathbf{z}$ ,  $\mathbf{z}^*$ , that is, matrices formed by the minors of order  $k$ . The operators  $(1 - q^2)^{-1/2} T(z_a^\alpha)$ ,  $(1 - q^2)^{-1/2} T((z_a^\alpha)^*)$ ,  $a = 1, 2, \dots, n$ ,  $\alpha = 1, 2, \dots, m$ , are respectively the  $q$ -analogues of creation and annihilation operators. The "creation operators" are placed in the right hand side of (5) to the left of the "annihilation operators". This allows one to produce a  $q$ -analogue of the polynomial  $\det(1 - \mathbf{z}\mathbf{z}^*)$  in a standard way as follows.

Let  $1 \leq \alpha_1 < \alpha_2 < \dots < \alpha_k \leq m$ ,  $1 \leq a_1 < a_2 < \dots < a_k \leq n$ . Introduce  $q$ -analogues of minors for the matrix  $\mathbf{z}$ :

$$\mathbf{z}^{\wedge k}_{\{a_1, a_2, \dots, a_k\}}^{\{\alpha_1, \alpha_2, \dots, \alpha_k\}} = \sum_{s \in S_k} (-q)^{l(s)} z_{a_1}^{\alpha_{s(1)}} z_{a_2}^{\alpha_{s(2)}} \dots z_{a_k}^{\alpha_{s(k)}},$$

with  $l(s) = \text{card}\{(i, j) \mid i < j \ \& \ s(i) > s(j)\}$  being the length of the permutation  $s$ .

The  $q$ -analogue  $y \in \text{Pol}(\text{Mat}_{mn})_q$  for the polynomial  $\det(1 - \mathbf{z}\mathbf{z}^*)$  is defined by

$$y = 1 + \sum_{k=1}^m (-1)^k \sum_{\{J' \mid \text{card}(J')=k\}} \sum_{\{J'' \mid \text{card}(J'')=k\}} \mathbf{z}^{\wedge k}_{J''} \cdot \left( \mathbf{z}^{\wedge k}_{J'} \right)^*.$$

Let  $\lambda > m + n - 1$ . Now (4) allows one to define the integral with weight  $y^\lambda$  as follows:

$$\int_{\mathbb{U}_q} f d\nu_\lambda \stackrel{\text{def}}{=} C(\lambda) \text{tr}(T(f)T(y)^\lambda q^{-2\Gamma(\tilde{\rho})}), \quad f \in D(\mathbb{U})_q,$$

with  $C(\lambda) = \prod_{j=0}^{n-1} \prod_{k=0}^{m-1} (1 - q^{2(\lambda+1-N)} q^{2(j+k)})$  being a normalizing multiple that provides

$$\int_{\mathbb{U}_q} 1 d\nu_\lambda = 1.$$

The Hilbert space  $L^2(d\nu_\lambda)_q$  is defined as a completion of the space  $D(\mathbb{U})_q$  of finite functions with respect to the norm  $\|f\|_\lambda = \left( \int_{\mathbb{U}_q} f^* f d\nu_\lambda \right)^{1/2}$ . The closure  $L_a^2(d\nu_\lambda)_q$  in  $L^2(d\nu_\lambda)_q$  of the unital subalgebra  $\mathbb{C}[\text{Mat}_{mn}]_q \subset \text{Pol}(\text{Mat}_{mn})_q$  generated by  $z_a^\alpha$ ,  $a = 1, 2, \dots, n$ ,  $\alpha = 1, 2, \dots, m$ , will be called a weighted Bergman space.

Note that the relations (1) and the algebra  $\mathbb{C}[\text{Mat}_{mn}]_q$  were considered in many works on quantum groups (see [2]). The  $*$ -algebra  $\text{Pol}(\text{Mat}_{mn})_q$  determined by the relations (1) and (2), is a  $q$ -analogue of the Weyl algebra. This becomes plausible after one performs the 'change of variables' as follows:

$$z_a^\alpha \mapsto (1 - q^2)^{-1/2} z_a^\alpha; \quad (z_a^\alpha)^* \mapsto (1 - q^2)^{-1/2} (z_a^\alpha)^*.$$

Consider the orthogonal projection  $P_\lambda$  in  $L^2(d\nu_\lambda)_q$  onto the weighted Bergman space  $L_a^2(d\nu_\lambda)_q$ . It is possible to show that  $P_\lambda$  could be written as an integral operator (see [8])

$$P_\lambda f = \int_{\mathbb{U}_q} K_\lambda(\mathbf{z}, \boldsymbol{\zeta}^*) f(\boldsymbol{\zeta}) d\nu_\lambda(\boldsymbol{\zeta}), \quad f \in D(\mathbb{U})_q. \quad (6)$$

Our intention is to introduce the algebra  $\mathbb{C}[[\text{Mat}_{mn} \times \overline{\text{Mat}}_{mn}]]_q$  of kernels of integral operators and to determine an explicit form of the weighted Bergman kernel  $K_\lambda \in \mathbb{C}[[\text{Mat}_{mn} \times \overline{\text{Mat}}_{mn}]]_q$  involved in (6).

Introduce the notation

$$\mathbb{k}_i = \sum_{\substack{J' \subset \{1, 2, \dots, m\} \\ \text{card}(J')=i}} \sum_{\substack{J'' \subset \{1, 2, \dots, n\} \\ \text{card}(J'')=i}} z^{\wedge i}_{J''} \otimes \left( z^{\wedge i}_{J'} \right)^*. \quad (7)$$

Let  $\mathbb{C}[\overline{\text{Mat}}_{mn}]_q \subset \text{Pol}(\text{Mat}_{mn})_q$  be the unital subalgebra generated by  $(z_a^\alpha)^*$ ,  $a = 1, 2, \dots, n$ ,  $\alpha = 1, 2, \dots, m$ , and  $\mathbb{C}[\text{Mat}_{mn}]_q^{\text{op}}$  the algebra which differs from  $\mathbb{C}[\text{Mat}_{mn}]_q$  by a replacement of its multiplication law to the opposite one (this replacement is motivated in [8]). The tensor product algebra  $\mathbb{C}[\text{Mat}_{mn}]_q^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{mn}]_q$  will be called an algebra of polynomial kernels. It is possible to show that in this algebra  $\mathbb{k}_i \mathbb{k}_j = \mathbb{k}_j \mathbb{k}_i$  for all  $i, j = 1, 2, \dots, m$ .

We follow [7, 6] in equipping  $\text{Pol}(\text{Mat}_{mn})_q$  with a  $\mathbb{Z}$ -grading:  $\deg(z_a^\alpha) = 1$ ,  $\deg((z_a^\alpha)^*) = -1$ ,  $a = 1, 2, \dots, n$ ,  $\alpha = 1, 2, \dots, m$ . In this context one has:

$$\mathbb{C}[\text{Mat}_{mn}]_q^{\text{op}} = \bigoplus_{i=0}^{\infty} \mathbb{C}[\text{Mat}_{mn}]_{q,i}^{\text{op}}, \quad \mathbb{C}[\overline{\text{Mat}}_{mn}]_q = \bigoplus_{j=0}^{\infty} \mathbb{C}[\overline{\text{Mat}}_{mn}]_{q,-j}$$

$$\mathbb{C}[\text{Mat}_{mn}]_q^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{mn}]_q = \bigoplus_{i,j=0}^{\infty} \mathbb{C}[\text{Mat}_{mn}]_{q,i}^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{mn}]_{q,-j} \quad (8)$$

The kernel algebra  $\mathbb{C}[[\text{Mat}_{mn} \times \overline{\text{Mat}}_{mn}]]_q$  will stand for a completion of  $\mathbb{C}[\text{Mat}_{mn}]_q^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{mn}]_q$  in the topology associated to the grading in (8). The kernel algebra is constituted by formal series  $\psi = \sum_{i,j=0}^{\infty} \psi_{ij}$ , with  $\psi_{ij} \in \mathbb{C}[\text{Mat}_{mn}]_{q,i}^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{mn}]_{q,-j}$ .

Our main result is the following formula for the weighted Bergman kernel:

$$K_{\lambda} = \prod_{j=0}^{\infty} \left( 1 + \sum_{i=1}^m (-q^{2(\lambda+j)})^i \mathbb{k}_i \right) \cdot \prod_{j=0}^{\infty} \left( 1 + \sum_{i=1}^m (-q^{2j})^i \mathbb{k}_i \right)^{-1} \quad (9)$$

with  $\mathbb{k}_i$  being the polynomial kernels (7). (The right hand side of (9) determines an element of  $\mathbb{C}[[\text{Mat}_{mn} \times \overline{\text{Mat}}_{mn}]]_q$  since  $\mathbb{k}_i \in \mathbb{C}[\text{Mat}_{mn}]_{q,i}^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{mn}]_{q,-i}$  for all  $i = 1, 2, \dots, m$ ).

In the special case  $m = n = 1$  we get a well known result [5]:

$$\begin{aligned} K_{\lambda} &= \prod_{j=0}^{\infty} (1 - q^{2(\lambda+j)} z \otimes \zeta^*) \cdot \left( \prod_{j=0}^{\infty} (1 - q^{2j} z \otimes \zeta^*) \right)^{-1} = \\ &= \sum_{i=0}^{\infty} \frac{(1 - q^{2\lambda})(1 - q^{2(\lambda+1)}) \dots (1 - q^{2(\lambda+i-1)})}{(1 - q^2)(1 - q^4) \dots (1 - q^{2i})} z^i \otimes \zeta^{*i}. \end{aligned}$$

Now passage to a limit as  $q \rightarrow 1$  and replacement of  $\otimes$  by a dot yields  $K_{\lambda} \xrightarrow{q \rightarrow 1} (1 - z\zeta^*)^{-\lambda}$ .

A  $q$ -analogue of an ordinary Bergman kernel for the matrix ball (see [3]) is derivable from (9) by a substitution  $\lambda = m + n$ :

$$K = \prod_{j=0}^{m+n-1} \left( 1 + \sum_{i=1}^m (-q^{2j})^i \mathbb{k}_i \right)^{-1} \xrightarrow{q \rightarrow 1} (\det(1 - \mathbf{z} \cdot \zeta^*))^{-(m+n)}.$$

## References

- [1] R. J. Baston, M. G. Eastwood. The Penrose Transform. Its Interaction with Representation Theory. Clarendon Press. Oxford, 1989.
- [2] V. Chari, A. Pressley. A Guide to Quantum Groups, Cambridge Univ. Press, 1995.
- [3] L. K. Hua. Harmonic Analysis of Functions of Several Complex Variables. Transl. of Math. Monog., Vol. 6. Amer. Math. Soc., 1963.
- [4] N. E. Hurt. Geometric Quantization in Action. D. Reidel Publishing Company, Dordrecht-Boston-London, 1983.
- [5] S. Klimec, A. Lesniewski, *A two-parameter quantum deformation of the unit disc*, J. Funct. Anal. **115**, (1993), 1 – 23.

- [6] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. Function theory in the quantum matrix ball: an invariant integral. E-print: math.QA/9803110, 5 p.p.
- [7] S. Sinel'shchikov, L. Vaksman. *On  $q$ -analogues of Bounded Symmetric Domains and Dolbeault Complexes*, Mathematical Physics, Analysis and Geometry, **1**, (1998), 75 – 100; E-print: q-alg/9703005.
- [8] L. L. Vaksman, *Intertwining operators and quantum homogeneous spaces*, Math. Phys. Anal. Geom., **1** (1994), No 3/4, 329 – 409, and E-print: q-alg/9511007.